

Critical visibility for N -particle Greenberger-Horne-Zeilinger correlations to violate local realism

Marek Żukowski and Dagomir Kaszlikowski
Instytut Fizyki Teoretycznej i Astrofizyki
Uniwersytet Gdański, PL-80-952 Gdańsk, Poland
 (February 1, 2008)

Abstract

A sequence of Bell inequalities for N -particle systems, which involve three settings of each of the local measuring apparatuses, is derived. For GHZ states, quantum mechanics violates these inequalities by factors exponentially growing with N . The threshold visibilities of the multiparticle sinusoidal interference fringes, for which local realistic theories are ruled out, decrease as $(2/3)^N$.

Greenberger-Horne-Zeilinger (GHZ) correlations [1] lead to a strikingly more direct refutation of the argument of Einstein Podolsky and Rosen (EPR), on the possibility of introducing elements of reality to complete quantum mechanics [2], than considerations involving only pairs of particles. The EPR ideas are based on the observation that for some systems quantum mechanics predicts perfect correlations of their properties. However, in the case of three or more particles, in the entangled GHZ state, such correlations cannot be consistently used to infer at a distance hidden properties of the particles. In contradistinction to the original two particle Bell theorem, the idea of EPR, to turn the exact predictions of quantum mechanics against the claim of its completeness, breaks down already at the stage of defining the elements of reality.

The reasoning of GHZ involved perfectly correlated particle systems. However, the actual data collected in a real laboratory would reveal less than perfect correlations, and the imperfections of the particle collection systems would leave many of the potential events undetected. Therefore the original GHZ reasoning cannot be ever tested in the laboratory, and one is forced to make some modifications (already, e.g., in [3]).

To face these difficulties several N -particle Bell inequalities appeared in the literature [4], [5]. All these works show that quantum predictions for GHZ states violate these inequalities by an amount that grows exponentially with N . The increasing number of particles, in this case, does not bring us closer to the classical realm, but rather, makes the discrepancies between the quantum and the classical more profound.

Any experimental realization of GHZ processes will be much more involved than a two-particle Bell test (for specific proposals see [6] and [7]). Therefore, the study of three or more particle interference effects does not seem to be a good route towards a loophole free test of the hypothesis of local hidden variables. However, other questions are to be answered in such experiments. For example, is it at all possible to observe three or four particle interferometric fringes which have no classical model? Before attempting such an experiment one must know the borderline between the quantum and the classical (i.e., local realism). According to current literature (with the exception of [5]) we enter the non-classical territory when the fringes in a N -particle interference experiment have visibilities higher than $2^{\frac{1}{2}(1-N)}$. The principal aim of this work is to show that, if one allows each of the local observers to have *three* measurements to choose from (instead of the usual *two*), the actual threshold is lower (for $N > 3$).

In a GHZ-Bell type experiment one has a source emitting N -particles each of which propagates towards one of N spatially separated measuring devices. The generic form of a GHZ N -particle state is

$$|\Psi(N)\rangle = \frac{1}{\sqrt{2}}(|+\rangle_1 \dots |+\rangle_N + |-\rangle_1 \dots |-\rangle_N). \quad (1)$$

Let us assume that the operation of each of the measuring apparatus is controlled by a knob which sets a parameter ϕ_l , and the l -th apparatus measures a dichotomic observable $O_l(\phi_l)$ with two eigenvalues ± 1 and the eigenstates defined by $|\pm, \phi_l\rangle_l = \frac{1}{\sqrt{2}}(|+\rangle_l \pm e^{i\phi_l}|-\rangle_l)$. The quantum prediction for obtaining specific results at the N measurement stations (for the idealized, perfect, experiment) reads

$$\begin{aligned} P_{QM}^{(N)}(r_1, r_2, \dots, r_N | \phi_1, \dots, \phi_N) \\ = \frac{1}{2^N} \left[1 + \prod_{l=1}^N r_l \cos \left(\sum_{k=1}^N \phi_k \right) \right], \end{aligned} \quad (2)$$

(r_l equal to -1 or $+1$). The GHZ correlation function is defined as

$$E^{(N)}(\phi_1, \dots, \phi_N) = \sum_{r_1, r_2, \dots, r_N=1} \prod_{l=1}^N r_l P^{(N)}(r_1, \dots, r_N | \phi_1, \dots, \phi_N), \quad (3)$$

and in the case of quantum mechanics, i.e. for $P^{(N)} = P_{QM}^{(N)}$, it reads $E_{QM}^{(N)} = \cos(\sum_{l=1}^N \phi_l)$.

From the perspective of local realism one can try to give a more complete specification of the state of a member of the ensemble of N -particle systems than the one given by $|\Psi(N)\rangle$. The usual approach is to define a space of hidden states Λ and a probability distribution $\rho(\lambda)$ of such states and to represent the probability of specific results by

$$P_{HV}^{(N)}(r_1, \dots, r_N | \phi_1, \dots, \phi_N) = \int_{\Lambda} d\lambda \rho(\lambda) \prod_{l=1}^N P_l(r_l | \lambda, \phi_l) \quad (4)$$

where $P_l(r_l | \lambda, \phi_l)$ is the probability to obtain result r_l in the l -th apparatus under the condition that the hidden state is λ and the macroscopic variable defining the locally measured observable is set to the value ϕ_l [8]. The locality of this description is guaranteed by independence of P_l on ϕ_i for all $i \neq l$.

We shall now derive a series of inequalities for the N -particle GHZ processes based on the following simple geometric observation [5]. Assume that one knows the components of a certain (Euclidean) vector $\vec{q}$ (the *known* vector), whereas about a second vector $\vec{h}$ (the *test* vector) one is only able to establish that its scalar product with $\vec{q}$ satisfies the inequality $\vec{h} \cdot \vec{q} < \|\vec{q}\|^2 = \vec{q} \cdot \vec{q}$. Then $\vec{h} \neq \vec{q}$. This simple geometric theorem can be extended to any pair of (real) objects for which one can define the scalar product (e.g., *matrices* or functions [5]).

We assume that each of the N spatially separated observers has three measurements to choose from. The local phases that they are allowed to set are $\phi_1^1 = \pi/6, \phi_2^1 = \pi/2, \phi_3^1 = 5\pi/6$ (for the first observer) and for all the other $N - 1$ observers they are $\phi_1^i = 0, \phi_2^i = \pi/3, \phi_3^i = 2\pi/3$, ($i = 2, \dots, N$).

Out of the quantum predictions for the N -particle correlation function at these settings one can construct a matrix endowed with N indices

$$E_{QM}^{(N)}(\phi_{i_1}^1, \dots, \phi_{i_N}^N) = \cos\left(\sum_{k=1}^N \phi_{i_k}^k\right) = Q_{i_1, \dots, i_N}^{(N)} \quad (5)$$

($i_k = 1, 2, 3$).

All that we know about local hidden variable theories is that their predictions (for the same set of settings as above) must have the following form:

$$E_{HV}^{(N)}(\phi_{i_1}^1, \dots, \phi_{i_N}^N) = \int d\lambda \rho(\lambda) \prod_{k=1}^N I_k(\lambda, \phi_{i_k}^k) = H_{i_1, \dots, i_N}^{(N)}, \quad (6)$$

where

$$I_k(\lambda, \phi_{i_k}^k) = \sum_{r_k} r_k P_k(r_k | \lambda, \phi_{i_k}^k). \quad (7)$$

Of course, in the case of a *deterministic* theory $I_k(\lambda, \phi) = \pm 1$ [8]. $H^{(N)}$ is our test matrix. Please note, that all one knows about $H^{(N)}$ is its structure.

The scalar product of two real matrices is defined by

$$(H^{(N)}, Q^{(N)}) = \sum_{i_1, \dots, i_N} H_{i_1, \dots, i_N}^{(N)} Q_{i_1, \dots, i_N}^{(N)}. \quad (8)$$

Our aim is to show the incompatibility of the local hidden variable description with the quantum prediction. To this end, we shall show that, for two or more particles,

$$\begin{aligned} & (Q^{(N)}, H^{(N)}) \\ & \leq 2^{N-1} \sqrt{3} < \|Q^{(N)}\|^2 = \frac{3^N}{2}. \end{aligned} \quad (9)$$

First, we show that $\|Q^{(N)}\|^2 = 3^N/2$. This can be reached in the following way:

$$\begin{aligned} \|Q^{(N)}\|^2 &= \sum_{i_1, \dots, i_N} \cos^2 \left(\sum_{k=1}^N \phi_{i_k}^k \right) \\ &= \frac{1}{2} \sum_{i_1, \dots, i_N} \left[1 + \cos \left(2 \sum_{k=1}^N \phi_{i_k}^k \right) \right] \\ &= \text{Re} \left\{ \sum_{i_1, \dots, i_N} \left[1 + \exp \left(2 \sum_{k=1}^N \phi_{i_k}^k \right) \right] \right\} \\ &= 3^N/2 + \text{Re} \left(\prod_{k=1}^N \sum_{i_k=1}^3 \exp(2i\phi_{i_k}^k) \right), \end{aligned} \quad (10)$$

where Re denotes the real part. Since $\sum_{l=1}^3 e^{i(l-1)(2/3)\pi} = 0$, the last term vanishes.

The scalar product $(H^{(N)}, Q^{(N)})$ is bounded from above by the maximal possible value of

$$S_\lambda^{(N)} = \sum_{i_1, \dots, i_N} \left[\cos \left(\sum_{k=1}^N \phi_{i_k}^k \right) \prod_{l=1}^N I_l(\lambda, \phi_{i_l}^l) \right], \quad (11)$$

and for $N \geq 2$

$$S_\lambda^{(N)} \leq 2^{N-1} \sqrt{3}. \quad (12)$$

To show (12), let us first notice that

$$S_\lambda^{(N)} = \text{Re} \left[\prod_{k=1}^N \sum_{i_k=1}^3 I_k(\lambda | \phi_{i_k}^k) \exp(i\phi_{i_k}^k) \right]. \quad (13)$$

For $k = 2, \dots, N$, one has $e^{i\phi_i^k} = e^{i[(l-1)/3]\pi}$ whereas for $k = 1$, $e^{i\phi_i^1} = e^{i(\pi/6)} e^{i[(l-1)/3]\pi}$. Thus, since $I(\lambda | \cdot) = \pm 1$, the possible values for

$$z_1^\lambda = \sum_{i_1=1}^3 I_1(\lambda | \phi_{i_1}^1) \exp(i\phi_{i_1}^1) \quad (14)$$

are $0, \pm 2e^{i\pi/2}, \pm 2e^{-i(\pi/6)}$, or finally $\pm 2e^{i(\pi/6)}$, whereas for $k = 2, \dots, N$ the possible values of

$$z_k^\lambda = \sum_{i_k=1}^3 I_k(\lambda | \phi_{i_k}^k) \exp(i\phi_{i_k}^k) \quad (15)$$

have their complex phases shifted by $\pi/6$ with respect to the previous set; i.e., they are 0, $\pm 2e^{i(2\pi/3)}$, ± 2 , or finally $\pm 2e^{i(\pi/3)}$. Since $|z_1^\lambda \prod_{k=2}^N z_k^\lambda| \leq 2^N$ and the minimal possible overall complex phase (modulo 2π) of $z_1^\lambda \prod_{k=2}^N z_k^\lambda$ is $\pi/6$, one has $\text{Re}(z_1^\lambda \prod_{k=2}^N z_k^\lambda) \leq 2^N \cos(\pi/6)$. Thus inequalities (12) and (9) hold.

The left inequality of (9) is a Bell inequality for the N -particle experiment. If one replaces $H^{(N)}$ by the quantum prediction $Q^{(N)}$ (compare (5)) the inequality is violated since

$$(Q^{(N)}, Q^{(N)}) = \frac{3^N}{2} > 2^{N-1}\sqrt{3}, \quad (16)$$

i.e., (9) is violated by the factor $(3/2)^N/\sqrt{3}$ (compare [4]).

The magnitude of violation of a Bell inequality is not a parameter which is directly observable in the experiment. It is rather the visibility of the N -particle interference fringes which can be directly observed. Further, the significance of all Bell-type experiments depends on the efficiency of the collection of the particles. Below a certain threshold value for this parameter experiments cannot be considered as tests of local realism. They may confirm the quantum predictions but are not falsifications of the hypothesis of local hidden variables. Therefore we will search for the critical visibility of N -particle fringes and collection efficiency, which do not allow anymore a local realistic model.

In a real experiment (under the assumption that quantum mechanics gives idealized, but correct predictions), the visibility of the N -particle fringes, $V(N)$, would certainly be less than 1. Also the probability of registering all potential events would be reduced by the overall collection efficiency. If one assumes that all N local apparata have the same collection efficiency η , and takes into account that these operate independently of each other, one can model the expected experimental results by

$$\begin{aligned} & P_{\text{expt}}^{(N)}(r_1, \dots, r_N | \phi^1, \dots, \phi^N) \\ &= \eta^N \left(\frac{1}{2}\right)^N \left(1 + V(N) \prod_{l=1}^N r_l \cos \sum_{k=1}^N \phi^k\right). \end{aligned} \quad (17)$$

The full set of events at a given measuring station consists now of the results $+1$ and -1 , when we succeed to measure the dichotomic observable, and a non-detection event (which is, in principle observable, if one uses event-ready state preparation [7]) for which one can introduce the value 0. The local realistic description requires that the probabilities of the possible events should be given by

$$\begin{aligned} & P_{\text{expt}}^{HV}(m_1, \dots, m_N | \phi^1, \dots, \phi^N) \\ &= \int d\lambda \rho(\lambda) \prod_{k=1}^N P_k(m_k | \lambda, \phi^k), \end{aligned} \quad (18)$$

with $m_i = +1, -1$ or 0. The local hidden variable correlation function for the experimental results (at the chosen settings) is now given by

$$E_{\text{expt } i_1, \dots, i_N}^{HV} = \int d\lambda \rho(\lambda) \prod_{k=1}^N I'_k(\lambda, \phi_{i_k}^k), \quad (19)$$

with

$$I'_k(\lambda, \phi_{i_k}^k) = \sum_{m_k=-1,0,+1} m_k P_k^{HV}(m_k | \lambda, \phi_{i_k}^k). \quad (20)$$

For deterministic models one has now $I'_k(\lambda, \phi_{i_k}^k) = 1, 0, -1$.

One can impose several symmetries on P_{expt}^{HV} . These symmetries are satisfied by the quantum prediction (17), and we can expect them to be satisfied in real experiments, within experimental error. The one that we impose here is that:

For all sets of results, $\{m_1, \dots, m_N\}$, that have equal number of zeros (one zero or more) the probability $P_{expt}^{HV}(m_1, \dots, m_N)$ has the same value, and this value is independent of the settings of the local parameters $\{\phi_{i_1}^1, \dots, \phi_{i_N}^N\}$.

One can define a function $f_N(m)$ which for $m = +1, -1, 0$ has the following values: $f(\pm 1) = \pm 1$, $f(0) = -1$ (compare [9]) and introduce auxiliary correlation function

$$\begin{aligned} \tilde{E}_{i_1, \dots, i_N} &= \int d\lambda \rho(\lambda) \sum_{m_1, \dots, m_N = -1, 0, +1} \\ &\times \prod_{k=1}^N [f(m_k) P_k(m_k | \lambda, \phi_{i_k}^k)] = \tilde{H}_{i_1, \dots, i_N}^{(N)}. \end{aligned} \quad (21)$$

Since, due to the symmetry conditions, one has, e.g., $\sum_{m_2=1, -1} f(m_2) P_{expt}^{HV}(0, m_2, m_3, \dots, m_N) = 0$, the following relation results:

$$\tilde{E}_{i_1, \dots, i_N} = E_{expt_{i_1, \dots, i_N}}^{HV} + [f(0)]^N P(0, \dots, 0), \quad (22)$$

where $P(0, \dots, 0)$ is the probability that all detectors would fail to register particles, and under our assumptions it is independent of the settings, and equals $(1 - \eta)^N$.

The auxiliary correlation function must satisfy the original inequality (9); i.e., one has

$$(Q^{(N)}, \tilde{H}^{(N)}) \leq 2^{N-1} \sqrt{3}. \quad (23)$$

However, this implies that

$$\begin{aligned} &-2^{N-1} \sqrt{3} - f(0)^N P(0, \dots, 0) q_{(N)} \\ &\leq (Q^{(N)}, E_{expt}^{HV}) \\ &\leq 2^{N-1} \sqrt{3} - f(0)^N P(0, \dots, 0) q_{(N)}, \end{aligned} \quad (24)$$

where

$$q_{(N)} = \sum_{i_1, \dots, i_N} Q_{i_1, \dots, i_N}^{(N)}. \quad (25)$$

Therefore, since if x is a possible value for $(Q^{(N)}, E_{expt}^{HV})$ then so is $-x$, one has

$$\begin{aligned} &|(Q^{(N)}, E_{expt}^{HV})| \\ &\leq 2^{N-1} \sqrt{3} - P(0, \dots, 0) |q_{(N)}|. \end{aligned} \quad (26)$$

Thus, we have obtained Bell inequalities of a form which is more suitable for the analysis of the experimental data.

The prediction (17) leads to the following correlation function

$$E_{expt}^{QM} = \eta^N V(N) E^{QM}, \quad (27)$$

which, when put into (26) in the place of E_{expt}^{HV} , gives the following relation between the critical visibility, $V_{cr}(N)$, and the critical collection efficiency, η_N^{cr} , for the N -particle experiment:

$$\eta_N^{crN} \frac{3^N}{2} V_{cr}(N) = 2^{N-1} \sqrt{3} - |q_{(N)}| (1 - \eta_N^{cr})^N. \quad (28)$$

The value of the expression $q_{(N)}$ can be found in the following way:

$$\begin{aligned} q_{(N)} &= \sum_{i_1, \dots, i_N} \cos \left(\sum_{k=1}^N \phi_{i_k} \right) \\ &= \text{Re} \left(\sum_{i_1, \dots, i_N} \prod_{k=1}^N \exp(i\phi_{i_k}^k) \right) \\ &= \text{Re} \left(\prod_{k=1}^N \sum_{i_k} \exp(i\phi_{i_k}^k) \right) \\ &= \text{Re} \left[2^N i \exp \left(i(N-1) \frac{\pi}{3} \right) \right] = -2^N \sin \left((N-1) \frac{\pi}{3} \right). \end{aligned} \quad (29)$$

The critical value of the visibility of the multiparticle fringes decreases now faster than in the earlier approaches [4]. For perfect collection efficiency, ($\eta = 1$), it has the lowest value, which is

$$V_{cr}(N) = \sqrt{3} \left(\frac{2}{3} \right)^N, \quad (30)$$

and, if $N \geq 4$, it is lower than $(\frac{1}{\sqrt{2}})^{N-1}$. The specific values for several particles are $V_{cr}(2) = 77.8\%$, $V_{cr}(3) = 51.3\%$, $V_{cr}(4) = 34.2\%$, $V_{cr}(5) = 22.8\%$ and $V_{cr}(10) = 3\%$, whereas the standard methods lead to $V(2)^{old} = 70.7\%$, $V(3)^{old} = 50.0\%$, $V(4)^{old} = 35.4\%$, $V(5)^{old} = 25.0\%$ and $V(10)^{old} = 4.4\%$. This suggests that for the original GHZ problem (four particles) one should rather aim at making experiments which allow for three settings at each local observation station. Surprisingly, the measurements should not be performed for the values for which we have perfect GHZ-EPR correlations (i.e the values for which the correlation function equals to ± 1) [10].

The critical efficiency of the particle collection also decreases with growing N , and for perfect visibilities it reads $\eta^{(2)} = 87.0\%$, $\eta^{(3)} = 79.8\%$, $\eta^{(4)} = 76.5\%$, $\eta^{(5)} = 74.4\%$. The gain over the inequalities [4] is in this respect very small, and begins again at $N = 4$. However, for very big N the critical efficiency is close to $\frac{2}{3}$ (compared with $\frac{1}{\sqrt{2}}$ for [4]).

M. Żukowski acknowledges support of the University of Gdańsk research grant no. BW-5400-5-0306-7, and of the 1996/97 Austrian-Polish Scientific-Technological Collaboration Program PRO22, and a Visiting Professorship of the University of Innsbruck.

REFERENCES

- [1] D.M. Greenberger, M.A. Horne and A. Zeilinger, in *Bell's theorem and the Conception of the Universe*, edited by M. Kafatos (Kluwer Academic, Dordrecht, 1989).
- [2] A. Einstein, B. Podolsky, and N. Rosen, Phys. Rev. **47** (1935) 777.
- [3] D.M. Greenberger, M. Horne, A. Shimony and A. Zeilinger, 1990, Am. J. Phys., **58**, 1131.
- [4] N.D. Mermin, Phys. Rev. Lett. **65**, 1838 (1990); S.M. Roy and V. Singh, Phys. Rev. Lett. **67**, 2761 (1991); M. Ardehali, Phys. Rev. A **46**, 5375 (1992); A.V. Belinskii and D.N. Klyshko, Physics-Uspekhi **36** 653 (1993). The inequalities involve pairs of settings at each of the local measuring apparata. Quantum mechanics violates these inequalities maximally by the factor $\sqrt{2^{N-1}}$.
- [5] M. Żukowski, Phys. Lett. A **177** 290 (1993); M. Żukowski, in *Quantum Interferometry*, eds. F. DeMartini, A. Zeilinger, (World Scientific, Singapore, 1994)
- [6] For example K. Wódkiewicz, L.-W. Wang and J.H. Eberly, Phys. Rev. A **47**, 3280 (1993), J.I. Cirac and P. Zoller, Phys. Rev. A **50**, R2799 (1994), S. Haroche, Ann. N. Y. Acad. Sci. **755**, 73 (1995), T. Sleator and H. Weinfurter, Ann. N. Y. Acad. Sci. **755**, 646 (1995), F. Laloë, Current Science **68**, 1026 (1995), C.C. Gerry, Phys. Rev. A **53**, 2857 (1996). T. Pfau, C. Kurtsiefer and J. Mlynek, Quant. and Semiclass. Opt. **8**, 665 (1996). Also, essentially all designs for quantum computer gates can produce GHZ states.
- [7] B. Yurke and D. Stoler, Phys. Rev. Lett. **68**, 1251 (1992); M. Żukowski, A. Zeilinger, M.A. Horne and A.K. Ekert, Phys. Rev. Lett. **71**, 4287 (1993); M. Żukowski, A. Zeilinger and H. Weinfurter, Ann. Phys. N. Y. Acad. Sci. **755**, 91 (1995); A. Zeilinger, M.A. Horne, H. Weinfurter and M. Żukowski, Phys. Rev. Lett., **78**, 3031 (1997).
- [8] We present our results only for the case of deterministic local hidden variable theories (for which $P_l(r_l|\lambda, \phi_l) = 1$, or 0), but the generalization to stochastic theories is obvious.
- [9] A. Garg and N.D. Mermin, Phys. Rev. D, **35**, 3638 (1987).
- [10] Seemingly, in the proposed inequalities altogether 3^4 global settings are involved, but as 27 elements the matrix (5), for $N = 4$, vanish, thus effectively, if one does not intend to make a delayed choice experiment, one may perform measurements at only 54 global settings.